ET: Astronomy 230 Section 1– MWF 1400-1450

134 Astronomy Building

This Class (Lecture 11):

Habitable Planets

Next Class:

Planets of Life

HW #3 is due Friday.

Presentation on Weds Carl Thomas Hassan Bhayani Aaron Bowling

Presentations Sept 26 Andrew Coughlin

Nicolas Jaramillo Chris Fischetti

Music: *Blister In the Sun* – The Violent Femmes

Sept 19, 2005

Astronomy 230 Fall 2004

Outline

- The formation of the Earth– atmosphere and oceans.
- What is n₂?
 - Breaks into 2 terms.
 - Whole lecture for n_e

presents his annual slideshow talk A Tour of the Night Sky

Professor

Emeritus **James Kaler**

Astronomer, author, teacher of generations of students.

Tuesday, September 20th,7 pm 119 Materials Science and Engineering Building, (NW corner of Green and Matthews)

Come and let Professor Kaler show us what we can see in the sky, and tell us about the beauty of the universe and our place in it.

The event is free and all are welcome (and no, you don't need to be an astronomer vourself to enjoy it!)



• Extra Credit Assignment

• Go to slideshow and write

1-2 paragraphs (typed!)

learned at the show and

about something you

1-2 paragraphs about

something that would

• Worth $\frac{1}{4}$ of a homework

impact Alien Life

(Astro 230).

assignment. • Due on Sept 26th.

L.W. Looney

Sept 19, 2005

Earth's Surface

Astronomy 230 Fall 2004

- 70% of the Earth's surface is covered with water
 - Ocean basins
 - Sea floors are young, none more than 200 million years old
- 30% is dry land Continents
 - Mixture of young rocks and old rocks
 - Up to 4.2 billion years old







L.W. Loonev

L.W. Looney

Geologically Active Surface

- The young rocks on the Earth's surface indicate it is geologically active
- Where do these rocks come from?
 - Volcanoes
 - Rift valleys
 - Oceanic ridges
- Air, water erode rocks
- The surface is constantly changing



Recycling Bio-elements

- From gravity and radioactivity, the core stays hot.
- This allows a persisting circulation of bioelements through continental drift— melting of the crust and re-release through volcanoes.
- Otherwise, certain elements might get locked into sediment layers- e.g. early sea life.
- Maybe planets being formed now, with less supernovae, would not have enough radioactivity to support continental drifts and volcanoes. (Idea of Peter Ward and Donald Brownlee.)



http://www.pahala-hawaii.com/j-page/image/activevolcanoe.jpg

```
Sept 19, 2005
```

L.W. Looney

```
Sept 19, 2005
```

Astronomy 230 Fall 2004

```
L.W. Looney
```

The Earth's 1st Atmosphere



- The interior heat of the Earth helped with the Earth's early atmosphere.
- The inner disk had most gases blown away and the proto-Earth was not massive enough to capture these gases. And any impacts (e.g. the moon), would have blown the atmosphere away.
- Most favored scenario is that comets impacted that released water (H₂O), carbon dioxide (CO₂), and Nitrogen (N₂)– the first atmosphere.
- The water condensed to form the oceans and much of the CO₂ was dissolved in the oceans and incorporated into sediments- such as calcium carbonate (CaCO₃).

Our Atmosphere

Astronomy 230 Fall 2004

Ì

- Rocks with ages greater than 2 million years show that there was little or probably no oxygen in the Earth's atmosphere.
- The current composition: 78% nitrogen, 21% oxygen, and trace amounts of water, carbon dioxide, etc.
- Where did the oxygen come from?
- Cyanobacteria made it.
 - Life on Earth modifies the Earth's atmosphere.



http://www.uweb.ucsb.edu/~rixfury/conclusion.htm

L.W. Looney

This New Planet

- Mostly oceans and some solid land (all volcanic).
- Frequent impacts of remaining planetesimals (ending about 3.8 billion years ago).
- Impacts would have sterilized the young Earth– Mass extinctions and maybe vaporized oceans (more comets?).
- Impacts and volcanic activity created the continental landmasses.
- Little oxygen means no ozone layerultraviolet light on the surface.
- Along with lightning, radioactivity, and geothermal heat, provided energy for chemical reactions.

Sept 19, 2005	
---------------	--

```
Astronomy 230 Fall 2004
```





Complex term, so let's break it into two terms:

- n_p: number of planets suitable for life per planetary system
- f_s: fraction of stars whose properties are suitable for life to develop on one of its planets

 $n_e = n_p \times f_s$



L.W. Looney



Water



- Water is a key to life on Earth.
- Primary constituent of life- "Ugly bags of mostly water"
 - Life is about 90% water by mass.
- Primary role as a solvent
 - Dissolves molecules to bring nutrients and remove wastes. Allows molecules to "move" freely in solution.
 - Must be in liquid form, requiring adequate pressure and certain range of temperatures.
- This sets a requirement on planets, if we assume that all life requires water.

Water as a Solvent

- The water molecule is "polar". The oxygen atoms have more build-up of negative charge than the hydrogen. This allows water molecules to link up, attracted to each other.
- In this way, water attracts other molecules, surrounds them and effectively dissolves them into solution.



L W LABORES

Sept 19, 2005

Astronomy 230 Fall 2004

Water

- A very good temperature buffer
 - Absorbs significant heat before its temperature changes
 - When it vaporizes, it takes heat with it, cooling down its original location
- It floats.
 - Good property for life in water.
 - Otherwise, a lake would freeze bottom up, killing life.
 - By floating to the surface, it can insulate the water somewhat.



Example: Dissolving Table Salt

The partial charges of the water molecule are attracted to the Na⁺ and Cl⁻ ions. The water molecules work their way into the crystal structure and between the individual ions, surrounding them and slowly dissolving the salt.



Keeping it Useful: Atmposphere 🚺

- Need to have enough pressure to keep water from boiling away at low temperature
 - Cooking at higher elevation requires more time.
 Boiling point lowered: water doesn't get as hot.
 - If pressure too low, water goes directly from ice to vapor (like dry ice CO₂)
- On the other hand, high pressure may make life more difficult to form.
- In addition, the range of temperature for Earth based complex life is less than 325K.

Astronomy 230 Fall 2004

Astronomy 230 Fall 2004

L.W. Looney

Keeping It Warm, but not too Warm

- What controls a planets temperature?
 - The amount of light received from its star.
 - The amount of energy the planet reflects back.
 - And any Greenhouse effects of the planet.
- Earth's greenhouse effect raises its temperature by about 15%.
- Given a star's luminosity, a range of acceptable temperatures translates into a range of distances to the star.
- This range is called the star's habitable zone (HZ), as planets in this range have temperatures suited for life.
- Only a rough guideline.

 Long living s 	ar		
• Planets with s temps)	table orbits (thus sta	able	Mars
 Liquid Water 			
Heavy Eleme	nts– C, N, O, etc.		Earth
Protection fro	m UV radiation		Mercury
0.	5M _{Sun} star		
	Astronomy 230 Fal	1 2004	I W Looney

Habitable Zones-

Galactic Habitable Zone

Astronomy 230 Fall 2004

• Likewise the galaxy has regions that are better suited to life.

Sept 19, 2005

- In the inner regions of our galaxy, supernovae are too frequent.
- In the outer regions, there are too few metals.
- Simulation of Galaxy Zone from early stages to now.



http://astronomy.swin.edu.au/GHZ/GHZmovie.html

The Sun's Variation

Ì

- As the Sun ages, it gets slightly brighter.
- When it was younger, its luminosity was 70% current values.
- A young Earth should have been 20K colder-iceball!
- During our ice ages, the temperature only changed by about 1%.



http://www.cherishclaire.com/iceball.htm

Astronomy 230 Fall 2004

L.W. Looney

L.W. Looney

L.W. Looney

The Sun's Variation

- There is evidence that the Earth did nearly freeze over-2.8 billion years ago and 700 million years ago.
- Probably changes in the Greenhouse gases.
- This implies that the habitable zone can vary with time, thus the real habitable zone is smaller than shown before
- Some have postulated that real zone is only 0.95 to 1.01 AU! If the Earth were 1% farther away- Iceballed. And n_p would be very small ~ 0.1.



http://www.soest.hawaii.edu/gerard/GG108/images/bylot.jpg

L.W. Loonev

Earth's Atmosphere

- Most recent studies suggest an efficient planet negativefeedback mechanism (like a thermostat).
- CO₂ cycles from atmosphere (greenhouse gas) and oceans (buried sedimentary especially carbonate rock).
- CO₂ in atmosphere: temporarily dissolved CO₂ in rainfall reacts with weathered rocks.
- Carbon is buried and can be released by volcanoes.
- Negative feedback process
 - Increase in temperature: evaporation of oceans, more rainfall, more weathering and CO_2 reduction, so decrease in temperature.
 - This negative feedback stabilizes the Earth's temperature.



```
Sept 19, 2005
```

Life Adds to Feedback

Astronomy 230 Fall 2004

• Life increases the weathering of rock.

Sept 19, 2005

- J.E. Lovelock has proposed that life also stabilizes the planet temperature.
- Regardless, the negative feedback helps with the habitable zone, so we can estimate perhaps n_p is more around 1- more Earth chauvinism?





Optimism?

Astronomy 230 Fall 2004

- Carl Sagan argues for $n_p > 3$.
 - If Venus had less clouds (less greenhouse) it could have been cool enough for life.
 - If Mars had a thicker atmosphere it could have been warm enough for life.
 - If solvents other than water were used. maybe the moons of the outer planets?
 - Giant Jupiter-like planets close in?

http://www.uranos.eu.org/biogr/sagane.html http://spider.ipac.caltech.edu/staff/jarrett/sagan/sagan.html







Pessimism?

- We only considered temperature. What about:
 - Gravity
 - Atmospheric pressure?
 - Size of the moon or planet?
 - Does life need a Moon-like moon? Does life need the tides? Does the Moon protect the Earth's rotation? Is a Jupiter needed?
- If we impose Earth chauvinism, we can easily reduce to $n_p \sim 0.1$



Sept 19, 2005

Astronomy 230 Fall 2004 L.W. Looney http://sagiru.tripod.com/Travel/Lost_in_the_Sahara/lost_in_the_sahara

 $n_e = n_p \times f_s$

- n_p: number of planets suitable for life per planetary system
- f_s: fraction of stars whose properties are suitable for life to develop on one of its planets
- We can list 5 situations that will have an effect on f_s.



n_p

- Can range from 0.01 to >3.
 - Is seismic activity necessary to recycle bioelements?
 - How important is the first atmosphere? Ozone?
 - Is a moon needed? A large Jupiter-like planet?
 - Is liquid water a requirement? Other solvents okay?
 - Not too hot, not too cold; not too much pressure, not too little.
 - Habitable Zone around the star.
 - Galactic Habitable Zone
 - Does atmosphere need feedback mechanism?
 - But in our solar system, maybe 5 nearly possible life planets.

Astronomy 230 Fall 2004

Sept 19, 2005

Differences of Stars to Life



L.W. Looney

1. <u>Metal rich stars</u>. Stars with heavy elements, probably more likely to have planets. Suggested in the current planet searches. About 90% of all stars have metals.



http://nike.cecs.csulb.edu/~kjljvioWallpaporsPlanet s%2001.jpg

Sept 19, 2005

Differences of Stars to Life

2. <u>Main sequence stars</u>. Need the brightness to stay as constant as possible. Otherwise the temperature changes dramatically on the planets. This is 99% of all stars.



Sept 19, 2005

L.W. Looney

Differences of Stars to Life

Astronomy 230 Fall 2004



<u>Binarity</u>. Planets may form. But they may have odd orbits unless the 2 stars are far enough apart or the planet orbits the pair. Only 30% of all stars are single stars. 50% of all stars are single stars or wide binary stars.



http://spaceflightnow.com/news/n0210/11planet/



Astronomy 230 Fall 2004

L.W. Looney

Secondary Star

Gamma Cephei System

Planet

Differences of Stars to Life



- **3.** <u>Length of time on the main sequence</u>. We need temperature stability for 5 x 10⁹ years to get intelligence on Earth. This rules out stars more massive than 1.25 solar masses! 90% of all stars are less massive than that.
- 4. <u>Minimum mass of star</u>. If ice exists close to the star, that would imply the formation of Jupiter-like planets not Earth-like planets. And, any life bearing planet would have to be closer to the star– and closer to stellar effects (e.g. tidal locking and more flares from low mass stars). That limits us to a minimum of 0.5 solar masses, about 25% of all stars.

```
Sept 19, 2005
```

Astronomy 230 Fall 2004

L.W. Looney

Adding it all up

	Stellar Requirement	Mass Limit	Fraction OK	Cumulative Fraction
✓	Heavy Elements		0.9	0.9
✓	Main Sequence		0.99	0.891
	Main Sequence Lifetime	M < 1.25 M _{sun}	0.90	
	Synchronous Rotation/ Flares	$M > 0.5 \ M_{Sun}$	0.25	
	Not a Binary		0.30	0.267
	Wide Binary Separation		0.50	

Sept 19, 2005

Astronomy 230 Fall 2004

Adding it all up

1	
Ŷ	
Д	

L.W. Looney

Stellar Requirement	Mass Limit	Fraction OK	Cumulative Fraction
Heavy Elements		0.9	0.9
Main Sequence		0.99	0.891
Main Sequence Lifetime	M < 1.25 M _{sun}	0.90	0.890
Synchronous Rotation/ Flares	$M > 0.5 M_{Sun}$	0.25	0.200
Not a Binary		0.30	
Wide Binary Separation		0.50	

f_s

Stellar Requirement	Mass Limit	Fraction OK	Cumulative Fraction
Heavy Elements		0.9	
Main Sequence		0.99	
Main Sequence Lifetime	M < 1.25 M _{sun}	0.90	
Synchronous Rotation/ Flares	$M > 0.5 \ M_{Sun}$	0.25	
Not a Binary		0.30	
Wide Binary Separation		0.50	

Sept 19, 2005

Astronomy 230 Fall 2004

Sept 19, 2005

Astronomy 230 Fall 2004

L.W. Looney