Astronomy 330

This class (Lecture 10):

Nature of Life
Allison Hanna
Nick Mucia

Next Class:

Origins of Life
Yi Sun
Cheryl Cwik

HW 4 is due Thursday

Music: *Life on Mars?* – David Bowie

Presentations

- **Allison Hanna**: Life on Mars
- **Nick Mucia**: Self-replicating Probes

Outline

- What is a nucleic acid?

Drake Equation

That's 0.67 Life-liking systems/year

Frank Drake

$$N = R_* \times f_p \times n_e \times f_l \times f_i \times f_c \times L$$

N	R_*	f_p	n_e	f_l	f_i	f_c	L
# of advanced civilizations we can contact in our Galaxy today	Star formation rate	Fraction of stars with planets	# of Earthlike planets per system	Fraction on which life arises	Fraction that evolve intelligence	Fraction that commun-icate	Lifetime of advanced civilizations
	19 stars/ yr	0.4 systems/ star	1.25 x 0.07 = 0.0875 planets/ system	life/ planet	intel./ life	comm./ intel.	yrs/ comm.

Nucleic Acid: DNA and RNA

- Two types of nucleic acid.
- A polymer built up from monomers we'll come back to which ones.
- RNA (RiboNucleic Acid) is usually a long strand
- DNA (DeoxyriboNucleic Acid) is the double helix– visualize as a spiral ladder.



Nucleic Acid: DNA and RNA

- These molecules carry the genetic information of the organism– the message that gets coded into the amino acid chain.
- It is very much like computer code in many ways– and teaches how to spell useful word (proteins) out of the letters of the available amino acids.



DNA / RNA

- The origins of DNA and RNA are mysterious and amazing
- DNA/RNA are complex: Built from three basic types of monomers
  1. Sugar (deoxyribose or ribose)
  2. A phosphate PO_4
  3. One of four "nitrogenous bases"
     - Adenine (A)
     - Guanine (G)
     - Cytosine (C)
     - Thymine (T) in DNA / Uracil (U) in RNA



Sugars: Ribose or Deoxyribose



We will represent the sugar molecule (either ribose or deoxyribose) as a pentagon with two eyes.

Sugars: Ribose or Deoxyribose





 http://www.dscc.edu/bwilliams/Biology/biology1molemodels.htm

Phosphates

- Is often referred to as phosphoric acid.
- Makes five bonds with oxygen.







Phosphates and Sugars

- Make the sides of the twisted DNA ladder structure.
- Sugars and phosphates connect up in alternating bonds. P-S-P-S-P-S-P
- These are phosphodiester bonds.



And the Bases

5 types in 2 groups (based on structure):

- Purines:
  - Adenine
  - Guanine
- Pyrimidines:
  - Cytosine
  - Uracil
  - Thymine



http://www.bmrb.wisc.edu/referenc/nomenclature/figures/bases.gif

For DNA

For RNA

Purines: Adenine

- 5-sided ring built on the side of a 6-sided ring.



Adenine (A)



A

Adenine

$H_5C_5N_5$

http://resources.emb.gov.hk/biology/english/inherit/genetics.html
http://dlm.tmu.edu.tw/phase2/glossary/image/adenine.gif

Purines: Guanine

- 5-sided ring built on the side of a 6-sided ring.



Guanine (G)



G

Guanine

$H_5C_5N_5O$

http://resources.emb.gov.hk/biology/english/inherit/genetics.html
http://dlm.tmu.edu.tw/phase2/glossary/image/adenine.gif

Pyrimidines: Cytosine

- 6 sided rings (without a 5 sided ring)



Cytosine (C)



C

Cytosine

$H_5C_4N_3O$

http://resources.emb.gov.hk/biology/english/inherit/genetics.html
http://dlm.tmu.edu.tw/phase2/glossary/image/adenine.gif

Pyrimidines: Uracil

- 6 sided rings (without a 5 sided ring)



Uracil (U)



U

Uracil

$H_4C_4N_2O_2$

For RNA

http://nautilus.fis.uc.pt/molecularium/stereo/
http://dlm.tmu.edu.tw/phase2/glossary/image/adenine.gif

Pyrimidines: Thymine

- 6 sided rings (without a 5 sided ring)



Thymine (T)



Thymine

$H_6C_5N_3O_2$

For DNA

http://resources.emb.gov.hk/biology/english/inherit/genetics.html

http://dlm.tmu.edu.tw/phase2/glossary/image/adenine.gif

Monomers and Polymers

Monomer:

1. Amino acids → Proteins
2. Sugar → Nucleic acids
   phosphate
   nitrogenous bases

Polymer:

1. Proteins
2. Nucleic acids

Making RNA Mean Something

- Schematic of an RNA molecule.
- This segment can be read from left to right as ACU– called a codon (a three letter word, so to speak)
- Can be translated to a specific amino acid (**the code!**) – this corresponds to the amino acid Threonine.
  GGU is gylcine.
- By building up these amino acid codons, we can spell out (and thus construct) a protein.



Meaning in Mystery

FIRST LETTER	SECOND LETTER U	SECOND LETTER C	SECOND LETTER A	SECOND LETTER G	THIRD LETTER
U	Phenylalanine	Serine	Tyrosine	Cysteine	U
	Phenylalanine	Serine	Tyrosine	Cysteine	C
	Leucine	Serine	Stop	Stop	A
	Leucine	Serine	Stop	Tryptophan	G
C	Leucine	Proline	Histidine	Arginine	U
	Leucine	Proline	Histidine	Arginine	C
	Leucine	Proline	Glutamine	Arginine	A
	Leucine	Proline	Glutamine	Arginine	G
A	Isoleucine	Threonine	Asparagine	Serine	U
	Isoleucine	Threonine	Asparagine	Serine	C
	Isoleucine	Threonine	Lysine	Arginine	A
	(Start) Methionine	Threonine	Lysine	Arginine	G
G	Valine	Alanine	Aspartate	Glycine	U
	Valine	Alanine	Aspartate	Glycine	C
	Valine	Alanine	Glutamate	Glycine	A
	Valine	Alanine	Glutamate	Glycine	G

For DNA replace U with T

http://library.thinkquest.org/C004535/PF_amino_acids.html

Overconstrained

- 4 options for each letter in the Codon
- 4 x 4 x 4 = 64 options (can think if it as bits)
- But only 20 amino acids ⇒ over constrained
- 4 x 4 = 16 wouldn't work.
- Life picked the next highest number and copes with redundancy.

FIRST LETTER	SECOND LETTER U	SECOND LETTER C	SECOND LETTER A	SECOND LETTER G	THIRD LETTER
U	Phenylalanine	Serine	Tyrosine	Cysteine	U
	Phenylalanine	Serine	Tyrosine	Cysteine	C
	Leucine	Serine	Stop	Stop	A
	Leucine	Serine	Stop	Tryptophan	G
C	Leucine	Proline	Histidine	Arginine	U
	Leucine	Proline	Histidine	Arginine	C
	Leucine	Proline	Glutamine	Arginine	A
	Leucine	Proline	Glutamine	Arginine	G
A	Isoleucine	Threonine	Asparagine	Serine	U
	Isoleucine	Threonine	Asparagine	Serine	C
	Isoleucine	Threonine	Lysine	Arginine	A
	(Start) Methionine	Threonine	Lysine	Arginine	G
G	Valine	Alanine	Aspartate	Glycine	U
	Valine	Alanine	Aspartate	Glycine	C
	Valine	Alanine	Glutamate	Glycine	A
	Valine	Alanine	Glutamate	Glycine	G

DNA

- For life more complicated than viruses, the genetic code is stored in DNA.
- Differs from RNA in a few ways: uses deoxyribose sugar rather than ribose sugar and it uses thymine instead of uracil.
- Forms the double strand where two complementary bonds are held together with weaker hydrogen bonding– allowing easier separation.
- In that case, bases form unique pairs:
  - AT, TA, GC, CG

http://mbsu.sus.mcgill.ca/POST_MIDTERM PICS/DNA is my life.jpg

DNA





- A codon of DNA: AT, CG, TA
  - purine to pyrimidine connections



The Double Helix

- Resembles a twisted ladder
- The sides of the DNA ladder are made of the sugar and phosphate.
- The steps or rungs of the ladder are composed of one of the 4 nitrogenous base pairs.
  - AT, TA, GC, CG
- In other words, if you know the sequence on one side, you can deduce the sequence on the other side.



The Double Helix

- The ladder is twisted into the helix shape since the hydrogen bonds are at an angle.
- 3 pairs make up a codon, like RNA (4x4x4 = 64)
- Each codon is info on the amino acid, but only 20 of those– again over constrained.





Genes

- Each codon specifies an amino acid, and a sequence of condons specifies a protein or enzyme.
- E. coli bacterium has about 4,000 genes, and at any time those genes specify about 1,000 enzymes. Many genes are duplicates.



Ta-Backy

- Different organisms have different number of genes.
- Tobacco mosaic virus has 4 genes.
- A small bacterium has about 1000 genes– average sized bacterium has 4000 genes.



TOBACCO MOSAIC VIRUS



http://pathmicro.med.sc.edu/mhunt/intro-vir.htm

My Old Blue Genes

- The Human Genome Project found 30,000 genes
- If you took all of the nucleic acid in one human cell and stretched out the long sequence, it would be more than a meter long!
- Human cells have 3 x 10^9 base pairs, but 98% of it has no obvious function, and 99.9% is the same for all humans.

http://images.encarta.msn.com/xrefmedia/sharemed/targets/images/pho/t373/T373681A.jpg

My Old Blue Genes

- This 98% is often called “junk” DNA, but it is still unclear what it’s function is..
- May control the early development from embryo to adult.
- May be as important as the protein encoding portion, but we don’t know.
- There is evidence that there is evolutionary conservation of "junk" DNA, which implies importance.

http://images.encarta.msn.com/xrefmedia/sharemed/targets/images/pho/t373/T373681A.jpg

Feb 14, 2008 Astronomy 330 Spring 2008

Chromosomes

- Best way to package DNA is in chromosomes– DNA wrapped around proteins,
- Humans have 23 pairs of chromosomes (total of 46).
- Each ranges from 50 million to 250 million base pairs
- For each set, you got half from each parent.



http://folding.stanford.edu/education/GAH/gene.html

Feb 14, 2008 Astronomy 330 Spring 2008

Which requires the most genes?

1. Onion
2. Mosquito
3. Carp
4. Human



http://www.thefishermom.com/images/071804small.htm
http://www.themoderatevoice.com/files/joe-mosquito.jpg
http://www.freewebs.com/flyingonion/Onion.gif

Feb 14, 2008 Astronomy 330 Spring 2008

Chromosomes



http://gslc.genetics.utah.edu/units/basics/tour/chromosome.swf

DNA: Message in a Cell

- A cell is informed it needs a enzyme– call it Z.
- Other enzymes in nucleus unravel and separate the easily broken DNA at the site where the gene for making that enzyme in encoded.



http://www.accessexcellence.org/AB/GG/mRNA.html

DNA: Message in a Cell

- Transcription of the gene is made via complementary bases and are assembled in a messenger RNA or mRNA.
- DNA zips itself back together.
- The mRNA (a series of codons) moves from the nucleus to the cytoplasm.



http://www.accessexcellence.org/AB/GG/mRNA.html

DNA: Message in a Cell

- Translation is the next step.
- A ribosome (the site of the protein synthesis) recognizes the mRNA by a special base sequence that attaches.
- The amino acids are built up from transfer RNA (tRNA) that move along the mRNA.
- The tRNAs have anticodon and carry amino acids.
- The chain of amino acids grows until the stop codon signals the completion of enzyme Z.



http://www.accessexcellence.org/AB/GG/mRNA.html

Reproduction

- DNA unzips itself, with appropriate enzyme.
- Each strand acts like a template for making a new strand.
- As each side is complementary, the molecule is successfully reproduced into 2 copies.

http://xupacabras.weblog.com.pt/arquivo/zipper.jpg

Reproduction

- For dividing cells, a copy goes to each daughter cell.
- Really, the process includes many special enzymes, so sometimes errors can occur.
- Still, very efficient
- DNA is the stuff from which all life is made.
- Probably not the method of the first life– too complicated.



http://xupacabras.weblog.com.pt/arquivo/zipper.jpg

Molecular Basis of Life

1. Atoms needed are H,O,N, and C with small amounts of P and S.
2. 2 basic molecules are essential for life: proteins and nucleic acids
3. Both are polymers– made of simpler monomers that make up the "alphabet" or code of life. These direct the transcription and translation of the proteins from the code.
4. Proteins and nucleic acids are closely linked at a fundamental level. Communicating through the genetic code that must have originated very early. In most cases, the same code is used by different messages for chicken or shark or human or even Elvis.

Molecular Basis of Life

5. #4 rises an important question.
   - Proteins synthesis must be directed by nucleic acids, but nucleic acid transcription requires enzymes (proteins).
   - Chicken or the egg problem?
   - Did proteins arise on Earth first and give rise to nucleic acids, or vice versa?

Chirality

Handedness: Some molecules exist in two versions based on the position of the bonds. One molecule is the mirror image of the other, but they are not similar.



Chirality



http://universe-review.ca/I15-32-chirality.jpg

We are Left-Handed

- Amino acids in non-biological situations are mixtures of both, but in life only left-handed molecules are used.
- Why? We don't know.



http://www.sp.uconn.edu/~bi107vc/fa02/terry/proteins.html

We are Left-Handed

- To match, sugars in life are right-handed
- Suggests a common ancestor for life.
- The opposite should have worked just as well, and this arrangement probably arose out of chance. Once a preponderance of one chirality occurred it was replicated

ET's Food Limitations

An ET organism may be made of the same stuff, but if they are made of right-handed amino acids, they couldn't eat our food. Bummer.





From Space?

- The Murchison meteorite (Australia 1969) contained approximately even amount of left and right amino acids
- 70 different amino acids were found in it, but only 6 are used in living organisms.
- New results show that 4 of the amino acids had a slight excess of left-handed types.

First Life

- We currently think that life appeared on Earth around 3.8 billion years ago, or only 700 million years after the formation of the Earth. (Based on fossil evidence)
- That is about the same time as the heavy bombardment ended. So, that means life was fast– perhaps only a few 10-100 million years from sterile planet to party town.



http://youconnect.canon-europe.com/swedish/2003-10/images/earth/love_parade.gif

Life

- The most crucial monomers required for life are:
  - Amino acids (20 flavors) for **proteins**
  - The nucleotides: sugar, phosphates, and nitrogenous bases for **DNA/RNA**.
- How did they occur in a useful configuration so **fast** on the early Earth?
  - Remember the early Earth is not a fun place.
    - Poisonous gas atmosphere, hot, lots of meteorites, and cable TV is still 3.8 billion years away.

Molecular Basis of Life

i. Two basic molecules are essential for life: proteins and nucleic acids
ii. Both are polymers– made of simpler monomers
iii. Proteins and nucleic acids are closely linked at a fundamental level.
iv. Did proteins arise on Earth first and give rise to nucleic acids, or vice versa? Or from space?
v. This leads us to the chemical evolution of life.

Chemical Evolution

- Chemical basis of life obviously crucial.
- Apparently evolution of life is a continuation of tendencies toward greater complexity
- Chemical evolution has 3 steps:
  - Synthesis of monomers
  - Synthesis of polymers from the monomers
  - Transition to life.