

Astronomy 330



This class (Lecture 16):
Biological Evolution

Next Class:

Biological Evolution
Brent Nelson
Ryan Ross

Music: *Spaceboy* – Smashing Pumpkins


March 11, 2008

Astronomy 330 Spring 2008

Midterm



- Highest scores ever..
- Average = 97.4
- Median = 99.5



March 11, 2008

Astronomy 330 Spring 2008

HW #3



- **Brent Nelson:**
<http://www.greatdreams.com/reptlan/reps.htm>
- **Mark Mazzone:**
<http://www.webpan.com/dsinclair/ufo.html>

March 11, 2008

Astronomy 330 Spring 2008

Outline



- Two types of cell life: Eukaryotes and Prokaryotes.
- All life can be divided into 3 types:
 - Bacteria
 - Archaea
 - Eukarya
- Genetic diversity of life...

March 11, 2008

Astronomy 330 Spring 2008

Drake Equation



Frank Drake

That's 2.9 life systems/decade



$$N = R_* \times f_p \times n_e \times f_l \times f_i \times f_c \times L$$

# of advanced civilizations we can contact in our Galaxy today	Star formation rate	Fraction of stars with planets	# of Earthlike planets per system	Fraction on which life arises	Fraction that evolve intelligence	Fraction that communicate	Lifetime of advanced civilizations
	19 stars/yr	0.4 systems/star	1.25 x 0.07 = 0.0875 planets/system	0.44 life/planet	intel./life	comm./intel.	yrs/comm.

March 11, 2008

Astronomy 330 Spring 2008



Evolution of Intelligence



- What is intelligence?
- “The ability to model the world, including the organism’s own self” is a workable definition.
 - A spectrum of ability
- Crucial development for the full spectrum of intelligence is the diversity of life on Earth.
- Intelligence is not a requirement of life.



<http://www.amonline.net.au/insects/images/site/insect1.jpg>

March 11, 2008

Astronomy 330 Spring 2008



Evolution of Intelligence



- First, we will examine the diversity of life; the fossil record shows a huge diversity with time.
- Organisms range from bacteria to humans.
- 1.8 x 10⁶ known species
 - Insects account for most (1.0 x 10⁶)
 - Estimated that only 10% are known.
 - Bacteria are hard to classify—only 7700 species so far.



<http://www.amonline.net.au/insects/images/site/insect1.jpg>

March 11, 2008

Astronomy 330 Spring 2008



Evolution of Intelligence



- Remember that all of these organisms use nearly identical genetic codes, so life descended from a common ancestor.
- Primary challenge of biology is to explain how life from a single type of organism, diversified so much.
- Evolution is the primary concept.



March 11, 2008

Astronomy 330 Spring 2008

Life



If we took all the biomass of all the animals, and all the biomass of all the viruses, bacteria, protozoa, and fungi— who weighs more?

Around 90% of all biomass on the Earth is in the smallest and simplest lifeforms.




March 11, 2008

Astronomy 330 Spring 2008

Bacteria



- 40 million bacterial cells in a gram of soil
- 1 million bacterial cells in a milliliter of fresh water
- Something like five nonillion (5×10^{30}) bacteria in the world.



Staph bacteria
http://www.scharfphoto.com/fine_art_prints/archives/000608.php

March 11, 2008

Astronomy 330 Spring 2008

You or not you?



- This is more non-you cells in your body than you-cells in your body!
 - You are outnumbered 10 to 1!
 - Mostly on your skin and in your digestive track



Bacteria under a toe-nail
<http://news.nationalgeographic.com/news/2007/02/070206-skin-microbes.html>


March 11, 2008

Astronomy 330 Spring 2008

Classification of Life



1. Prokaryotes
 - No cell nucleus— DNA floating around
 - Always single-cell creatures like bacterium
 - Came first
 - Outnumber and outweigh the second class (eukaryotes)



Copyright © 2004 Pearson Education, Inc., publishing as Benjamin Cummings.

March 11, 2008


Astronomy 330 Spring 2008

Classification of Life



2. Eukaryotes

- Have a cell nucleus, a membrane to protect the DNA
- Basis of all multi-cell creatures
- Also some single-cell creatures like amoebas.
- DNA arranged into chromosomes in nucleus- 23 pairs for humans.



March 11, 2008

Astronomy 330 Spring 2008

Prokaryotes




Divided into 2 domains:

1. Eubacteria or “true” bacteria
 - 20% of the world’s biomass.
 - Thought to be the oldest surviving organisms.
 - Often found in harsh environments: hot springs, undersea vents, salty seashores, etc, which were probably more common on the early Earth.
 - Some evidence that ancient organisms were heat-lovers (maybe)
2. Archaea

Prokaryotes

- **Smaller**
- **Simpler**



- **Most do not have membrane-enclosed organelles**
- **Bacteria and archaea**

Carl Woese here at UIUC, discovered Archaea scheme.

Copyright © 2004 Pearson Education, Inc., publishing as Benjamin Cummings.

March 11, 2008

Astronomy 330 Spring 2008


Eukaryotes



- All animals, plants, and fungi.

Eukaryotes

- **Larger**
- **More complex**



- **Membrane-enclosed organelles**
- **Protists, plants, fungi, animals**


March 11, 2008

Astronomy 330 Spring 2008

3 Domains of Life



- Genetically speaking, Archaea and Eukarya are more similar to one another than are Bacteria and Archaea
- Implies that Archaea and Bacteria split and then all Eukarya split from Archaea
- A major implication for the evolution of life on Earth



The old “kingdom” classification is no longer really used, such as plant kingdom or animal kingdom


March 11, 2008

Astronomy 330 Spring 2008

Genetic Relations



- This is a major change from the old methods of assigning groups based on outward form and anatomy.
- Instead based on studies of the genetic code.
- Surprise: Human and chimpanzees share about 99% of the same DNA, and about 97% with mice.
- Surprise: 2 species of fruit fly look very much alike, but only share about 25%. Some of this differences is due to “junk” DNA.




<http://www.uglybug.org/index00.shtml>

<http://www.pritchettcartoons.com/fruitfly.htm>

March 11, 2008

Astronomy 330 Spring 2008

Changes in Bio-Systems



- Today’s view: evolution is the most important and unifying property of life.
- Anaximander (c. 610–547 BC): life arose in water and gradually became more complex
- Empedocles (c. 492–432 BC): survival of the fittest (but, “a good idea stated within an insufficient theoretical frame loses its explanatory power and is forgotten” by Hans Reichenbach)
- Aristotle (384–322 BC): species are fixed and independent of each other → evolution discarded for 2000 years
- Fossil record: slowly broke down the Aristotelian theory

March 11, 2008

Astronomy 330 Spring 2008

For the Species Survival



1 Population with varied inherited traits



2 Elimination of individuals with certain traits



3 Reproduction of survivors



4 Increasing frequency of traits that enhance survival and reproductive success

- Darwin (1809–1882) & Malthus (1766-1834):
 - Populations can grow faster than food sources can support them.
 - Creates a struggle for survival that can wipe out competitors.
 - Individual variations has advantages or disadvantages in the struggle for survival
 - Natural selection can create unequal reproductive success

Copyright © 2001 by Benjamin Cummings, an imprint of Addison Wesley.


March 11, 2008

Astronomy 330 Spring 2008

Filling the Niche with Finch



- Other Evidence:
 - Adapted species in the Galápagos Islands, in particular finches
 - Artificial breeding of house/farm animals and vegetables
- DNA is really the mechanism of natural selection, but evolution requires both heredity and environment



March 11, 2008

Astronomy 330 Spring 2008

Mutant Sex

- Mutations from changes in the bases of DNA.
- Usually copying errors, but also radiation–radioactivity, cosmic rays, chemical agents, or UV light.
- About 3 mutations per person per generation.
- Most mutations are neutral, changes in the *junk* DNA.
- Why is sex important to this class?



http://www.mutantx.net/features/press_vwSexy.html

March 11, 2008

Astronomy 330 Spring 2008



Mutant Sex

- Sexual reproduction leads to greater genetic diversity– a difference between prokaryotes and eukaryotes?
- Asexual reproduction does not allow 2 new and beneficial mutations to combine.
- Blackberries have not changed much in 10 millions years, but sexual plants have produced: raspberries, thimbleberries, cloudberries, dewberries, etc.
- Sex is useful in the process, but the mutations are still key.



<http://www.alcasoft.com/arkansas/blackberry.html>

March 11, 2008


Astronomy 330 Spring 2008

Does it take a long time?



Cabbage, kale, kohlrabi, brussels sprouts, cauliflower and broccoli have same common ancestor– wild mustard. All bred by humans on a very short time scale.

This is selective breeding, but still the potential is in the DNA.



Or domestic lap dogs from wolves in about 5000 years.

pring 2008

Comparing Ages



- Important to understand history of Earth life is the ability to age different components
- Can be difficult
- Radioactive dating....
 - ^{14}C for the last 60,000 years
 - ^{40}K and ^{235}U for 100's of millions of years


March 11, 2008

Astronomy 330 Spring 2008


Radioactive Dating



Recall:



- Most atomic nuclei stable
- But some nuclei are *unstable*,
 \Rightarrow decay to new nucleus
“radioactive”



March 11, 2008

The Law of Radioactive Decay



As radioactive “parent” decays, the number of decay product or “daughters” increases

Decay Rule

Start out with N parents, 0 daughters

Time t since start	# parents	# daughters
0	N	0
$t_{1/2}$	$\frac{1}{2} N = \text{half as much}$	$\frac{1}{2} N$ have appeared
$2t_{1/2}$	$\frac{1}{4} N = \text{half again as much}$	$\frac{3}{4} N$
$3t_{1/2}$	$\frac{1}{8} N$	$\frac{7}{8} N$
$30t_{1/2}$	About $N/10^9$	99.9999999% N

Decay is a good “clock”

- Each radioactive species has different “tick”
- Rate= “half-life”
- Exponential decay from original population of n_0


March 11, 2008

Astronomy 330 Spring 2008

Radioactive Decay Examples




http://www.colorado.edu/physics/2000/isotopes/radioactive_decay3.html



March 11, 2008

Astronomy 330 Spring 2008

Radioactive Dating



Example 1: Carbon C=6p

- Carbon-12: 6p+6n, stable
- Carbon-14: 6p + 8n, unstable (1/2 life of 5730 years)
- $^{14}\text{C} \rightarrow ^{14}\text{N}$ (nitrogen)
- Nitrogen-14: 7p + 7n, stable

Example 2: Uranium U=92p

- Uranium-238: 92 p + 146 n (1/2 life of 4.5 billion years)


$^{238}\text{U} \rightarrow \text{chain of decays} \rightarrow ^{206}\text{Pb}$ (lead)

March 11, 2008

Carbon-14



- Cosmic rays from space are constantly hitting the Earth.
- React with ^{14}N in atmosphere to create ^{14}C .
- Decays back to ^{14}N with half life of 5730 years.
- But, there is an equilibrium in abundance
- In atmosphere, the ^{14}C is mostly in $^{14}\text{CO}_2$.



http://bbspot.com/Images/News_Features/2003/12/half-life.jpg

March 11, 2008

Astronomy 330 Spring 2008

Carbon-14



- Plants take in $^{14}\text{CO}_2$ with the $^{12}\text{CO}_2$ and other animals eat the plants.
- So, every living creature has a equilibrium ratio of $^{14}\text{CO}_2/^{12}\text{CO}_2$.
- When the organism dies, the ^{14}C decays to ^{14}N . By measuring how much ^{14}C remains, you can date the fossil.
- This works well to about 60,000 years.
 - Viking remains in Newfoundland– 500 yrs before Columbus.
 - Shroud of Turin to 1330 AD



<http://web.mit.edu/smeguire/www/newfoundland/newf16.html>


March 11, 2008

Astronomy 330 Spring 2008

Dating Rocks



- First you ask them out?
- No, you need a radioactive decay that has a longer half-life than ^{14}C .
- Potassium-argon
 - ^{40}K decays to ^{40}Ar with a 1200 Myr half-life.
- Uranium-lead
 - ^{235}U to ^{207}Pb with 700 Myr half-life.
- But these only work with volcanic layers.
- So, the ages of fossils are interpolated from ages of volcanic layers above and below them.



March 11, 2008

Astronomy 330 Spring 2008

<i>Era</i>	<i>Period</i>	<i>Myr Ago</i>	<i>Life Forms</i>	<i>Events</i>
Cenozoic	Quaternary	2	H. Sapiens	Ice ages
	Tertiary	65	Primates	Extinction of Dinosaurs
Mesozoic	Cretaceous	136	Birds	S. Atlantic open to 1900 miles
	Jurassic	190		N. Atlantic open to 600 miles
	Triassic	225	Mammals	Continental drift
Paleozoic	Permian	280	Reptiles	Pangaea breaks up
	Carboniferous	345	Amphibians	Formation of coal
	Devonian	395	Insects	
	Silurian	430	Land Plants	
	Ordovician	500	Fish	
Precambrian	Cambrian	543	Trilobites	
		545	Small Shelly Fossils	
		580	Ediacarans	
		600-800	Multicellular life	Snowball Earth episodes

March 11, 2008

Astronomy 330 Spring 2008

Increase of Complexity




- Last table showed only the last 800 Myrs.
- The more complex and intelligent organisms appeared towards the end.
- For many years it was thought that life originated in the Cambrian era, then Precambrian fossils were found.
- Then, it was realized that there were single-celled fossils that required microscopes.

<i>Myr Ago</i>	<i>Era</i>	<i>Event</i>
Now	Cenozoic	
	Mesozoic	
	Paleozoic	Macroscopic life/Snowball Earth
	Precambrian	
1000		Worm tracks
		Multicellular algae
		Eukaryotes certain
		Sexual reproduction
2000		Eukaryotes possible
	Protozoic	Oxygen-rich atmosphere
		Snowball Earth
		Formation of continents
3000	Archean	Life begins?
4000		Formation of Oceans
		Bombardment decreases
		Frequent impacts
	Hadean	Earth formed

Concepts



- As prokaryotes are simpler than eukaryotes, we expect them to exist first.
- Identifying fossil prokaryotes is difficult: they're tiny!
- But there is enough evidence that before 1500-2000 Myrs ago there are only prokaryotes fossils.
- Note: the oldest fossils (3800 Myrs ago) are under some dispute, but the 2800 Myr old fossils are universally accepted.
- All of the macroscopic life only arose in the last 600 Myrs– 1/6th of the history of life on Earth.



<http://www.earth.ox.ac.uk/research/geobiology/geobiology.htm>

Making Oxygen!

- The early prokaryotes played a crucial role for life on Earth by producing oxygen through photosynthesis.
- Cyanobacteria (also called blue-green algae) changed the world!
- Lived in colonies that formed mats or films, growing into large structures called stromatolites.
- Still around, but much more common before 700 Myrs ago.



Making Oxygen!



- Oxygen was new and important step in intelligence
- It allowed a new energy extraction method
 - Aerobic (using oxygen) metabolism
 - More complex life.
 - Created ozone layer (dry land now an option for life on Earth).



March 11, 2008

Astronomy 330 Spring 2008

Relationship to ETs



- Would evolution on other planets have a similar time-scale?
- Evolution is not a deterministic process.
- Selection seems to be mostly luck, rather than adaptation.
- On the other hand, many traits have developed in several lineages– warm blood and eyes.
- Some say that intelligence seems to increase in many lineages, so it is likely that if life exists then intelligent life exists.
- On the other hand, the plant kingdom never developed neurons.

March 11, 2008

Astronomy 330 Spring 2008

Summary



- This following slides are from:
<http://www.udayton.edu/~INSS/>
- Nice timeline of life on Earth.

March 11, 2008

Astronomy 330 Spring 2008